



Lilydale Clinic

484 Maroondah Highway
Lilydale VIC 3140
P (03) 9739 5244 F (03) 9735 5509

Yarra Glen Clinic

28 Bell Street Yarra Glen VIC 3775
P (03) 9730 1569

www.yarrarangesvet.com.au
clinic@yarrarangesvet.com.au

CYSTITIS AND LOWER URINARY TRACT DISEASE

What is cystitis?

Cystitis is a general term referring to inflammation in the urinary bladder. In cats, diseases of the lower urinary tract (the bladder and urethra) are grouped under the term 'feline lower urinary tract disease' or 'feline idiopathic cystitis' as in some cases there is no identifiable underlying cause.

What are the signs of cystitis?

Typical signs in cats with cystitis are those of inflammation and irritation of the lower urinary tract. The common signs are therefore:-

- Increased frequency and urgency of urination.
- Difficulty in urinating (spending a long time straining on the litter tray while passing only small quantities of urine).
- The presence of blood in the urine or blood spots in the litter tray.
- Occasionally complete obstruction to the passage of urine (straining persistently without producing any urine).

With the latter sign particularly (straining without the passage of any urine), it is important to seek urgent veterinary attention as complete blockage to the flow of urine can be a life-threatening complication if left untreated.

What causes feline cystitis?

There are a vast number of potential causes, but in many cats there may be severe inflammation of the bladder and/or urethra without an identifiable underlying cause (so called 'idiopathic'cystitis). These idiopathic cases have to be differentiated from other potential causes of the clinical signs though so that appropriate treatment can be given. Some of the potential causes are listed below:

- Idiopathic (inflammation for no known cause) but possibly caused by unrecognised viruses or bacteria
- Urinary calculi ('bladder stones') or crystals
- Bacterial infections
- Neoplasia (tumour)
- Anatomical abnormalities
- Urethral plugs (blockage with a mix of crystals, calculi/stones and/ or mucus)

How is cystitis diagnosed?

The initial diagnosis of is based on the identification of signs of lower urinary tract inflammation. If there there is doubt, analysis of a urine sample will confirm the presence of inflammation or blood.

Further investigation is usually required to attempt to identify the underlying cause of the cystitis.

What further investigations are required to diagnose the cause of FLUTD?

Where clinical signs are persistent or recurrent, a number of investigations may be required to differentiate idiopathic FLUTD from the other known causes of urinary tract inflammation.

These investigations may include:

- Laboratory analysis of a urine sample to look for inflammation, bacteria or crystals
- Bacterial culture of a urine sample
- Blood samples to look for other evidence of urinary tract disease or other systemic disease
- Radiographs (x-rays) and/or ultrasound examination of the bladder and urethra



Lilydale Clinic

484 Maroondah Highway
Lilydale VIC 3140
P (03) 9739 5244 F (03) 9735 5509

Yarra Glen Clinic

28 Bell Street Yarra Glen VIC 3775
P (03) 9730 1569

www.yarrarangesvet.com.au
clinic@yarrarangesvet.com.au

What is the treatment for cystitis?

This depends on the underlying cause. For example:

- Cases of idiopathic disease may respond to treatment with anti-inflammatory or analgesic ('pain-relieving') drugs, but it is crucial that you *only* use drugs specifically prescribed by your veterinary surgeon, as many human products are extremely dangerous to use in cats.
- Bacterial infections of the lower urinary tract, although uncommon in cats, will usually respond well to antibiotic therapy.
- If a cat develops a blocked urethra (this almost exclusively occurs in males), emergency treatment is required to remove the blockage, which may require flushing of the urethra while the cat is given a short anaesthetic.
- If bladder crystals are present (the most common are "struvite" crystals) they are usually able to be dissolved by using a special diet such as Hills Prescription diet s/d.

How can idiopathic cystitis be prevented?

Idiopathic cystitis is more common in cats that have a lower water consumption, and in cats that are inactive and obese. All these factors may relate, at least in part, to the frequency with which a cat urinates. Avoidance of obesity and encouraging exercise may be of some help and as cats tend to drink very little the feeding of at least some tinned food rather than exclusively a dry cat food product will help to maintain a higher water intake.